


Inferential Statistics and Probability a Holistic Approach

Chapter 2 Descriptive Statistics



This Course Material by Maurice Geraghty is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License.
Conditions for use are shown here: <https://creativecommons.org/licenses/by-sa/4.0/>

1

Measures of Central Tendency

- Mean
 - Arithmetic Average $\bar{X} = \frac{\sum X_i}{n}$
- Median
 - "Middle" Value after ranking data
 - Not affected by "outliers"
- Mode
 - Most Occurring Value
 - Useful for non-numeric data

2

Example

Anthony's Pizza, a Detroit based company, offers pizza delivery to its customers. A driver for Anthony's Pizza will often make several deliveries on a single delivery run. A sample of 5 delivery runs by a driver showed that the total number of pizzas delivered on each run

2 2 5 9 12

What is the Average?



- a) 2
- b) 5
- c) 6


3

Example – 5 Recent Home Sales

- \$500,000
- \$600,000
- \$600,000
- \$700,000
- \$2,600,000

4





- ### Measures of Variability
- Range
 - Variance
 - Standard Deviation
 - Interquartile Range (percentiles)
- 8

Range

$$\text{Max}(X_i) - \text{Min}(X_i)$$

$$125 - 67 = 58$$


9

Sample Variance

$$s^2 = \frac{\sum (x_i - \bar{x})^2}{n-1}$$

$$s^2 = \frac{\sum x_i^2 - (\sum x_i)^2 / n}{n-1}$$

10



Variance and Standard Deviation


X_i	$X_i - \bar{X}$	$(X_i - \bar{X})^2$	
2	-4	16	$s^2 = \frac{78}{4} = 19.5$
2	-4	16	
5	-1	1	
9	3	9	$s = \sqrt{19.5} \approx 4.42$
<u>12</u>	<u>6</u>	<u>36</u>	
30	0	78	

12

Interpreting the Standard Deviation

- Chebyshev's Rule
 - At least $100 \times (1 - (1/k)^2)\%$ of any data set must be within k standard deviations of the mean.
- Empirical Rule (68-95-99 rule)
 - Bell shaped data
 - 68% within 1 standard deviation of mean
 - 95% within 2 standard deviations of mean
 - 99.7% within 3 standard deviations of mean

13



Measures of Relative Standing

- Z-score
- Percentile
- Quartiles
- Box Plots

15

Z-score

- The number of Standard Deviations from the Mean
- $Z > 0$, X_i is greater than mean
- $Z < 0$, X_i is less than mean

$$Z = \frac{X_i - \bar{X}}{s}$$

16

Percentile Rank

Formula for ungrouped data

- The location is $(n+1)p$ (interpolated or rounded)
- n = sample size
- p = percentile

17

Quartiles

- 25th percentile is 1st quartile
- 50th percentile is median
- 75th percentile is 3rd quartile
- 75th percentile – 25th percentile is called the Interquartile Range which represents the "middle 50%"

18

IQR example

$n+1=31$

$.25 \times 31 = 7.75$ location 8 = **87** ← 1st Quartile

$.75 \times 31 = 23.25$ location 23 = **108** ← 3rd Quartile


Interquartile Range (IQR) = $108 - 87 = 21$

19

Box Plots

- A **box plot** is a graphical display, based on quartiles, that helps to picture a set of data.
- Five pieces of data are needed to construct a box plot:
 - Minimum Value
 - First Quartile
 - Median
 - Third Quartile
 - Maximum Value.

20



Outliers

- An outlier is data point that is far removed from the other entries in the data set.
- Outliers could be
 - Mistakes made in recording data
 - Data that don't belong in population
 - True rare events


22

Outliers have a dramatic effect on some statistics

- Example quarterly home sales for 10 realtors:

	2	2	3	4	5	5	6	6	7	50
	with outlier					without outlier				
Mean	9.00					4.44				
Median	5.00					5.00				
Std Dev	14.51					1.81				
IQR	3.00					3.50				

23



Using Z-score to detect outliers

- Calculate the mean and standard deviation without the suspected outlier.
- Calculate the Z-score of the suspected outlier.
- If the Z-score is more than 3 or less than -3, that data point is a probable outlier.

$$Z = \frac{50 - 4.4}{1.81} = 25.2$$

25

Outliers – what to do

- Remove or not remove, there is no clear answer.
- For some populations, outliers don't dramatically change the overall statistical analysis. Example: the tallest person in the world will not dramatically change the mean height of 10000 people.
- However, for some populations, a single outlier will have a dramatic effect on statistical analysis (called "**Black Swan**" by Nicholas Taleb) and inferential statistics may be invalid in analyzing these populations. Example: the richest person in the world will dramatically change the mean wealth of 10000 people.

26

Bivariate Data



- Ordered numeric pairs (X,Y)
- Both values are numeric
- Paired by a common characteristic
- Graph as Scatterplot

27

Example of Bivariate Data

- Housing Data
 - X = Square Footage
 - Y = Price

28



Correlation Analysis





- **Correlation Analysis:** A group of statistical techniques used to measure the strength of the relationship (correlation) between two variables.
- **Scatter Diagram:** A chart that portrays the relationship between the two variables of interest.
- **Dependent Variable:** The variable that is being predicted or estimated. "Effect"
- **Independent Variable:** The variable that provides the basis for estimation. It is the predictor variable. "Cause?" (Maybe!)


31

The Coefficient of Correlation, r

- **The Coefficient of Correlation (r)** is a measure of the **strength** of the relationship between two variables.
 - It requires interval or ratio-scaled data (variables).
 - It can range from -1 to 1.
 - Values of -1 or 1 indicate perfect and strong correlation.
 - Values close to 0 indicate weak correlation.
 - Negative values indicate an inverse relationship and positive values indicate a direct relationship.

32





- ### Causation
- Correlation does not necessarily imply causation.
 - There are 4 possibilities if X and Y are correlated:
 1. X causes Y
 2. Y causes X
 3. X and Y are caused by something else.
 4. Confounding - The effect of X and Y are hopelessly mixed up with other variables.
- 38

- ### Causation - Examples
- City with more police per capita have more crime per capita.
 - As Ice cream sales go up, shark attacks go up.
 - People with a cold who take a cough medicine feel better after some rest.
- 39

Formula for correlation coefficient r

$$r = \frac{SSXY}{\sqrt{SSX \cdot SSY}}$$

$$SSX = \sum X^2 - \frac{1}{n}(\sum X)^2$$

$$SSY = \sum Y^2 - \frac{1}{n}(\sum Y)^2$$

$$SSXY = \sum XY - \frac{1}{n}(\sum X \cdot \sum Y)$$

40


Example

- X = Average Annual Rainfall (Inches)
- Y = Average Sale of Sunglasses/1000
- Make a Scatter Diagram
- Find the correlation coefficient

X	10	15	20	30	40
Y	40	35	25	25	15

41

- ### Example *continued*
- Make a Scatter Diagram
 - Find the correlation coefficient
- 42



Example *continued*

X	Y	X ²	Y ²	XY
10	40	100	1600	400
15	35	225	1225	525
20	25	400	625	500
30	25	900	625	750
40	15	1600	225	600
115	140	3225	4300	2775

- $SSX = 3225 - 115^2/5 = 580$
- $SSY = 4300 - 140^2/5 = 380$
- $SSXY = 2775 - (115)(140)/5 = -445$

44

Example *continued*

$$r = \frac{SSXY}{\sqrt{SSX \cdot SSY}}$$

$$r = \frac{-445}{\sqrt{580 \cdot 380}} = -0.9479$$

- Strong negative correlation

45