


## Inferential Statistics and Probability a Holistic Approach

### Chapter 4 Probability

  
This Course Material by Maurice Geraghty is licensed under a Creative Commons Attribution-ShareAlike 4.0 International License. Conditions for use are shown here: <https://creativecommons.org/licenses/by-sa/4.0/>

1

## Probability

- Classical probability
  - Based on mathematical formulas
- Empirical probability
  - Based on the relative frequencies of historical data.
- Subjective probability
  - "one-shot" educated guess.

2

## Examples of Probability

- What is the probability of rolling a four on a 6-sided die?
- What percentage of De Anza students live in Cupertino?
- What is the chance that the Golden State Warriors will be NBA champions in 2018?

3

## Classical Probability

- Event
  - A result of an experiment
- Outcome
  - A result of the experiment that cannot be broken down into smaller events
- Sample Space
  - The set of all possible outcomes
- Probability Event Occurs
  - # of elements in Event / # Elements in Sample Space
- Example – flip two coins, find the probability of exactly 1 head.
  - {HH, HT, TH, TT}
  - $P(1 \text{ head}) = 2/4 = .5$

4

## Empirical Probability

- Historical Data
- Relative Frequencies
- Example: What is the chance someone rates their community as good or better?
  - $0.51 + 0.32 = 0.83$


**National: Rate Your community**

Rating	Percentage of Sample
Excel	32
Good	51
Fair	13
Poor	3
Other	1

5

## Rule of Complement


- Complement of an event
- The event does not occur
- $A'$  is the complement of  $A$
- $P(A) + P(A') = 1$
- $P(A) = 1 - P(A')$



6

### Additive Rule

- The **UNION** of two events A and B is that either A or B occur (or both). (All colored parts)
- The **INTERSECTION** of two events A and B is that both A and B will occur. (Purple Part only)
- Additive Rule:  
 $P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$



7


### Example

- In a group of students, 40% are taking Math, 20% are taking History.
- 10% of students are taking both Math and History.
- Find the Probability of a Student taking either Math or History or both.
- $P(M \text{ or } H) = 40\% + 20\% - 10\% = 50\%$

8

### Mutually Exclusive

- Mutually Exclusive
- Both cannot occur
- If A and B are mutually exclusive, then
  - $P(A \text{ or } B) = P(A) + P(B)$
- Example roll a die
  - A: Roll 2 or less    B: Roll 5 or more
  - $P(A)=2/6$      $P(B)=2/6$
  - $P(A \text{ or } B) = P(A) + P(B) = 4/6$



9

### Conditional Probability

- The probability of an event occurring GIVEN another event has already occurred.
- $P(A|B) = P(A \text{ and } B) / P(B)$
- Example: Of all cell phone users in the US, 15% have a smart phone with AT&T. 25% of all cell phone users use AT&T. Given a selected cell phone user has AT&T, find the probability the user also has a smart phone.
- A=AT&T subscriber    B=Smart Phone User
- $P(A \text{ and } B) = .15$      $P(A)=.25$
- $P(B|A) = .15/.25 = .60$

10

### Contingency Tables

- Two data items can be displayed in a contingency table.
- Example: auto accident during year and DUI of driver.

	Accident	No Accident	Total
DUI	70	130	200
Non- DUI	30	770	800
Total	100	900	1000

11

### Contingency Tables

	Accident	No Accident	Total
DUI	70	130	200
Non- DUI	30	770	800
Total	100	900	1000

Given the Driver is DUI, find the Probability of an Accident.

A=Accident    D=DUI

$P(A \text{ and } D) = .07$      $P(D) = .2$

$P(A|D) = .07/.2 = .35$

12

## Marginal, Joint and Conditional Probability

- **Marginal Probability** means the probability of a single event occurring.
- **Joint Probability** means the probability of the union or intersection of multiple events occurring.
- **Conditional Probability** means the probability of an event occurring given that another event has already occurred.

13


## Creating Contingency Tables

- You can create a hypothetical contingency table from reported cross tabulated data.
- First choose a convenient sample size (called a radix) like 10000.
- Then apply the reported marginal probabilities to the radix of one of the variables.
- Then apply the reported conditional probabilities to the total values of one of the other variable.
- Complete the table with arithmetic.

14

## Example

Create a two-way table from the cross tabulation of gender from the 2016 election results (from CNN)



15

## Example

First select a radix (sample size) of 10000

VOTED FOR	GENDER		Total
	Female	Male	
Trump			
Clinton			
Other			
Total			10000

16

## Example

Then apply the marginal probabilities to the radix (53% female, 47% male)

VOTED FOR	GENDER		Total
	Female	Male	
Trump			
Clinton			
Other			
Total	5300	4700	10000

17

## Example

Then apply the cross tabulated percentages for each gender. Make sure the numbers add up.

VOTED FOR	GENDER		Total
	Female	Male	
Trump	2173	2444	
Clinton	2862	1927	
Other	265	329	
Total	5300	4700	10000

18

### Example

Finally, complete the table using arithmetic.

VOTED FOR	GENDER		Total
	Female	Male	
Trump	2173	2444	4617
Clinton	2862	1927	4789
Other	265	329	594
Total	5300	4700	10000

19

### Multiplicative Rule

- $P(A \text{ and } B) = P(A) \times P(B|A)$
- $P(A \text{ and } B) = P(B) \times P(A|B)$
- Example: A box contains 4 green balls and 3 red balls. Two balls are drawn. Find the probability of choosing two red balls.
- A=Red Ball on 1<sup>st</sup> draw B=Red Ball on 2<sup>nd</sup> Draw
- $P(A)=3/7$   $P(B|A)=2/6$
- $P(A \text{ and } B) = (3/7)(2/6) = 1/7$

20

### Multiplicative Rule – Tree Diagram

21

### Independence

- If A is not dependent on B, then they are **INDEPENDENT** events, and the following statements are true:
  - $P(A|B)=P(A)$
  - $P(B|A)=P(B)$
  - $P(A \text{ and } B) = P(A) \times P(B)$

22

### Example

	Accident	No Accident	Total
DUI	70	130	200
Non- DUI	30	770	800
Total	100	900	1000

A: Accident      D:DUI Driver

$P(A) = .10$      $P(A|D) = .35 (70/200)$

Therefore A and D are **DEPENDENT** events as  $P(A) < P(A|D)$

23

### Example

	Accident	No Accident	Total
Domestic Car	60	540	600
Import Car	40	360	400
Total	100	900	1000

A: Accident      D:Domestic Car

$P(A) = .10$      $P(A|D) = .10 (60/600)$

Therefore A and D are **INDEPENDENT** events as  $P(A) = P(A|D)$

Also  $P(A \text{ and } D) = P(A) \times P(D) = (.1)(.6) = .06$

24

### Random Sample

- A **random sample** is where each member of the population has an equally likely chance of being chosen, and each member of the sample is **INDEPENDENT** of all other sampled data.

25

### Tree Diagram method

- Alternative Method of showing probability
- Example: Flip Three Coins
- Example: A Circuit has three switches. If at least two of the switches function, the Circuit will succeed. Each switch has a 10% failure rate if all are operating, and a 20% failure rate if one switch has already failed. Find the probability the circuit will succeed.

26

### Circuit Problem

27

### Switching the Conditionality

- Often there are questions where you desire to change the conditionality from one variable to the other variable
- First construct a tree diagram.
- Second, create a Contingency Table using a convenient radix (sample size)
- From the Contingency table it is easy to calculate all conditional probabilities.

28


### Example

- 10% of prisoners in a Canadian prison are HIV positive.
- A test will correctly detect HIV 95% of the time, but will incorrectly "detect" HIV in non-infected prisoners 15% of the time (false positive).
- If a randomly selected prisoner tests positive, find the probability the prisoner is HIV+

29

### Example

30



### Example

	HIV+ A	HIV- A'	Total
Test+ B	950	1350	2300
Test- B'	50	7650	7700
Total	1000	9000	10000

$$P(A | B) = \frac{950}{2300} \approx .413$$

31